

TUNISIE

HAMMAMET

du 19 | nov.
au 21 | 2021

4^e édition

AFRAMED 2021

VIH, Hépatites, Santé sexuelle
Infections émergentes

Dépistage et traitement de l'hépatite C en Méditerranée

Nabil DEBZI

Service Hépatologie
CHU Mustapha –Alger


www.aframed2021.org






Conflits d'intérêts

Speaker : Biotest, Gilead , Boston scientific , Roche




Oteros Jose (2014) Modelización del ciclo fenológico reproductor del olivo (Tesis Doctoral). Universidad de Córdoba, Córdoba,



Michaël Sofia

THE NOBEL PRIZE IN PHYSIOLOGY OR MEDICINE 2020



Illustrations: Niklas Elmehed

Harvey J. Alter


Michael Houghton

Charles M. Rice

“for the discovery of Hepatitis C virus”



Prévalence des patients virémiques et génotypage





Stratégies de Dépistage

- Dépistage institutionnel
- Micro-élimination
- Dépistage universel = Macro-Élimination



WHO Vision: Eliminate Viral Hepatitis as a Major Health Threat by 2030



**World Health
Organization**

“A world where viral hepatitis transmission is halted and everyone living with hepatitis has access to safe, affordable and effective care and treatment services”

**90% reduction in
new chronic HCV
infections**

**Treatment of 80% of
eligible persons with
chronic HCV infection**

**65% reduction in
mortality rates**



We still have a long way to go to achieve the WHO HCV elimination targets






Table 3 Comparative HBV and HCV screening policies in the non-EU Mediterranean countries

	Algeria	Egypt	Israel	Jordan	Lebanon	Libya	Morocco	Tunisia
Antenatal screening					HBV	HBV		HBV
Blood and organ donors	Both	Both	Both	Both	Both	Both	Both	Both
Blood transfusion or products prior to 1992 in EU, or any transfusion outside EU			Both	Both				
Clinical signs or laboratory signs (including cirrhosis and HCC)	Both	Both	HCV	Both	Both	Both		Both
Candidates for chemotherapy or immunosuppressive treatment			HBV		HBV	HBV		Both
Haemophiliacs who received concentration factors prior to 1987			Both		Both			
Haemodialysis	Both	Both	Both	Both	Both	Both	HBV	Both
History of shared injecting equipment	Both	Both	Both	Both	Both	Both		
History of long-term imprisonment	Both							
Hospital surgery patients						Both		
Household contacts	Both		Both	Both		Both		
HIV	Both				Both			
IVF candidates			Both			Both		
Men who have sex with men								
Migrants from high prevalence countries			Both					
Military recruits	Both	Both		Both	Both			Both
Organ or tissue transplants prior to 1992 in EU or outside EU			Both		Both			
Pre-employment		Both	HCV, health care	Both				
Pregnant women and newborns			Selective risk groups: HBV	HBV	HBV	HBV		HBV
Prenuptial	Both				HBV			HBV
STI clinic patients			Both	HBV	Both	HCV		
Traditional medicine exposure	HBV							
Unvaccinated healthcare workers	Both		HBV	Both				HBV
Occupational exposure and/or carrying out exposure-prone procedures	Both	Both	Both	Both	Both	HBV		HBV





Table 4 HCV screening recommendations included in the HCV Hellenic National Plan

Target population	HCV screening recommendations
General	Birth cohort screening: Adults born from 1945 to 1980
High-risk groups	<ul style="list-style-type: none"> • Persons with elevated transaminases • People who inject drugs (current and former IV drug users) • Recipients of a transfusion of blood, blood components, or an organ transplant before 1992 • Persons who are receiving or have received hemodialysis • Persons who have been parenterally exposed to potentially HCV infected medical instruments or paramedical procedures • Long-term steady sex partners of HCV-positive persons • Persons with a history of multiple sex partners • Children born to HCV-positive women • Persons with HIV infection • Persons with HBV infection • Incarcerated persons • Immigrants from high HCV prevalence countries

HBV, hepatitis B virus; HCV, hepatitis C virus



Patients with advanced liver disease
Haemophilia patients
Prisoners
Children
Patients engaged with drug treatment units
Migrant communities from high prevalence regions
People who inject drugs in networks
Men who have sex with men
Generational cohorts of high prevalence
Geographically defined areas





Prévalence VHC : 15.4 → 13.2%
 Nouveaux cas : MSM 1.9 → 3.9%
 Indication DAA : 11.4 → 61.5%
 2018 41.2% MSM
 Who Target : OUI sauf Nouvelles infections



HCV Micro-elimination in easy-to-screen population: decompensated cirrhosis

- HEPATHER : prospective cohort including 699 patients prior decompensation of HCV cirrhosis with a median follow-up of 37 months
- SVR : 86 % (84 % in patients with CP-C and/or MELD > 20)




Mortality: adjusted HR 0.44, 95% CI 0.26-0.74 p=0.002

- 571 DAA+ and 128 (18.3%) untreated patients (DAA-)



Mode	Key topics or content
Booklets	Four booklets in Arabic to raise awareness distributed in public places in the village—eg, mosques, schools, and health-care units Entitled: <i>How to protect yourself from catching hepatitis, How to live with hepatitis, How to support a patient with hepatitis, and Infection control measures for paramedical workers</i>
Animated cartoon films	Nine episodes of an Arabic 3D animated cartoon film titled <i>Abo Eloraif</i> (meaning "Father of Knowledge") each 2–3 min long; and regular broadcasting of <i>Abo Eloraif</i> through local channels Episodes covered modes of transmission, risk behaviours for transmission, and recommended behaviours to reduce transmission and social stigma
Song	"How a patient with viral hepatitis can protect his family and neighbours?"; taught to school children and circulated by use of public health broadcasts on local satellite channel Emphasis was put on the following points for the general public: how personal items (eg, sponge, towel, shaving tools, and nail cutter or scissors) should be personal to them, while at home, travelling, or at the barber shop or hairdressers; cups and needles for hijama (cupping) or acupuncture should be personal; and for health workers: any wound should be disinfected and covered; they should be committed to complying with infection control standards and use of non-reusable syringes; the importance of using sterile instruments for dental procedures, and doing renal dialysis and surgical procedures in a clean sterilised environment
Posters	Five posters in community and health-care facilities Key messages included prevalence of hepatitis worldwide and in Egypt; practices that do and do not transmit infection; people at higher risk and how to protect yourself from acquiring hepatitis C infection

Table 1: Educational materials used for awareness raising and public education¹⁴



Gamal Shiha et AL An educate, test, and treat programme towards elimination of hepatitis C infection in Egypt: a community-based demonstration project www.thelancet.com/gastrohep Published online July 17, 2018



“De-centralize” screening and care in communities

Patients may face difficulties in accessing testing and treatment facilities. Different populations have differing needs and require specific settings and measures in place to access treatment

Centralized



Treatment delivered through a bottleneck delays treatment initiation and risks losing the chance to connect patients to care


De-centralized



Decentralizing treatment allows streamlined access to care



HCV reinfection over 5 years







FIGURE 3 Key success factors for HCV elimination, a conceptual framework. HCV, hepatitis C virus



Deuffic-Burban S et al. Assessing the cost-effectiveness of hepatitis C screening strategies in France. J Hepatol (2018), <https://doi.org/10.1016/j.jhep.2018.05.027>



TABLE 3 Direct medical costs and health effects, by scenario, 2018-2031

Scenario	Cost (€ millions), 2018-2031	QALYs gained, 2018-2031	ICER relative to status quo (€/QALY)	ICER relative to previous least costly scenario (€/QALY)
Status quo	5463	-	-	
GHSS 2020 Targets 2023	Graduated screening 1	1968-1987 144 000	3552	3552
	Graduated screening 2	1948-1967 125 000	4532	^a
	Screening 1948-1977	142 000	4349	^a
	Screening 1958-1977	128 000	4831	^a
	Universal screening	145 000	6758	562 855


Note: Values have been rounded, so ICERs may not be reproducible using table values.

Abbreviations: ICER, incremental cost-effectiveness ratio; QALY, quality-adjusted life year; GHSS: Global Health Sector Strategy.

^aStrongly dominated scenario (costlier and less effective than another scenario).



Global timing of hepatitis C virus elimination in high-income countries




● High income mediterranean countries



Traitements

Traitements princeps

Traitements génériques





Recommended Treatment Regimens

- **Genotype-specific**

- Elbasvir/Grazoprevir: GT 1, 4
- Ledipasvir/Sofosbuvir: GT 1, 4, 5, 6


- **Pangenotypic**

- Sofosbuvir/Velpatasvir – GT 1-6
- Glecaprevir/Pibrentasvir – GT 1-6
- Sofosbuvir/Velpatasvir/Voxilaprevir – GT 1-6 (reserved for salvage therapy)




Pangenotypic Regimens: SOF/VEL for 12 Wks

ASTRAL-1^{[1]*}: SOF/VEL for 12 Wks in GT 1, 2, 4, 5, 6




ASTRAL-3^{†[2]}: SOF/VEL
for 12 Wks in GT 3





Pangenotypic Regimens: GLE/PIB for 8 Wks in Patients Without Cirrhosis



*Includes treatment-naïve and treatment-experienced patients.



HCV DAAs: Prices in USA and India versus Target

Drug	Current US price (lowest)	Current lowest Indian market price	Target price
Sofosbuvir	\$49,680	\$324	\$62
Daclatasvir	\$50,653	\$153	\$14
SOF+LDV	\$56,700	\$507	\$96
SOF+VEL	\$74,760	-	\$181-216

Gotham D, Barber M, Fortunak J, Pozniak A, Hill A. Rapidly falling costs for new hepatitis C direct-acting antivirals (DAAs): potential for universal access. Abstract number A-792-0516-01639, presented at AIDS2016, Durban.



Bioequivalent pharmacokinetics for generic and originator Hepatitis C Direct Acting Antivirals

Andrew M. Hill¹, Loai Tahat², Mohammed Khalil Mohammed³, Sanjay Nath⁴, Rabab Fayez Tayyem³, James A. Freeman⁵, Isma-hane Benbitour⁷, Sherine Helmy⁶; ¹Department of Translational Medicine, University of Liverpool, Liverpool, United Kingdom; ²Pharmaceutical Research Unit, Amman, Jordan; ³ACDIMA BioCentre, Amman, Jordan; ⁴Faculty of Medicine, Imperial College London, London, United Kingdom; ⁵GP2U Telehealth, Hobart, TAS, Australia; ⁶R&D Project Innovations, Pharco, Cairo, Egypt; ⁷BEKER Laboratories, Algiers, Algeria



AASLD 2017

DRUG	TRIAL TYPE	COMPANY	NUMBER	C _{MAX}	AUC _{0-∞}
Sofosbuvir	Four-way, four-period, fully replicated, single oral dose	EEPI	36	101.0 (88.1-115.7)	103.0 (97.6-109.7)
Daclatasvir	Two-way, two-period, single oral dose	Dawood	35	106.9 (100.2-114.0)	103.7 (98.3-109.4)
Sofosbuvir	Three-period, two-treatment, three sequence, semi-replicate	Beker	35	95.4 (84.7-107.5)	98.5 (91.6-106.0)
Daclatasvir	Three-period, two-treatment, three sequence, semi-replicate	Beker	35	104.1 (93.1-116.3)	103.0 (94.4-112.4)
Sofosbuvir	Three-period, two treatment, three sequence, partial replicate	Hetero	54	95.7 (87.2-105.2)	100.8 (96.2-105.6)
Sofosbuvir	-	Natco	-	96.1 (81.0-114.0)	100.7 (94.2-107.8)
Daclatasvir	-	Natco	-	94.5 (83.1-107.4)	96.5 (87.1-106.8)
Sofosbuvir	Two-period, two-treatment, single dose	Virchow	22	94.8 (83.3 – 107.9)	95.8 (86.9 – 105.7)



High SVR rates using parallel imported generic DAAs in 1160 patients with Hepatitis C

James Freeman; Nabil Debzi; Giten Khwairakpam; Julia Dragunova; Sergey Golovin; James Wang; Andrew Hill; Vicky Houghton-Price; Rachel Smith; Roxanna Korologou-Linden; John Freeman; Greg Jefferys.



World Hepatitis Summit 2017


SÃO PAULO, BRAZIL 1-3 NOVEMBER



Contents lists available at [ScienceDirect](https://www.sciencedirect.com)

Arab Journal of Gastroenterology

journal homepage: www.elsevier.com/locate/ajg



Editorial

Hepatitis C elimination in Africa: Seizing the moment for hepatitis-C free future

A. Cordie ,R. Mohamed , Mark W. Sonderup ,C. W Spearman , M. A. Medhat , N. Debzi , H. Desalegn , G. Esmat

Hepatitis Can't Wait

Delaying screening, diagnosis and initiation of DAAs will impact on achieving the WHO 2030 elimination goals.



Table 1. Participation in Screening and HCV Seroprevalence According to Sex.*

Variable	Men	Women	Total
Screening target population — no.†	32,207,165	30,298,399	62,505,564
Participated in screening — no. (%)‡	24,018,428 (74.57)	25,611,891 (84.53)	49,630,319 (79.40)
Previously treated for HCV infection with direct-acting antivirals since 2014 — no. (%)§	692,632 (2.88)	591,739 (2.31)	1,284,371 (2.59)
Screened for HCV antibodies — no. (%)¶	23,325,796 (97.12)	25,020,152 (97.69)	48,345,948 (97.41)
HCV seropositive			
No. of adults	1,252,443	976,885	2,229,328
Percent (95% CI)¶¶	5.37 (5.36–5.38)	3.90 (3.90–3.91)	4.61 (4.61–4.62)


* CI denotes confidence interval, and HCV hepatitis C virus.

† The screening target population included all Egyptian adults 18 years of age or older who had ever been issued a national identification number.

‡ Participation in screening was voluntary. Percentages are relative to the screening target population.

§ Percentages are relative to the number who participated in screening. Persons who had been previously treated with direct-acting antiviral agents were not screened for HCV antibodies.

¶ Percentages are relative to the number screened for HCV antibodies.



I.Waked et al, Special report Screening and Treatment Program to Eliminate Hepatitis C in Egypt
n engl j med 382;12 nejm.org March 19, 2020



Table 3. Cost of the Screening and Treatment Program.*

Variable	Value
Screening	
Staff cost — \$	
Medical teams	36,552,528
Administrative staff	15,000
Total	36,567,528
Medical supplies — \$	
Rapid diagnostic test for HCV antibodies	27,345,901
Consumables (e.g., gloves, swabs, and staff uniforms)	3,701,062
Total	31,046,963
Information technology and administration — \$	4,364,830
Overhead — \$	18,787,415
Total cost of HCV screening program — \$	90,766,736
No. with HCV seropositivity	2,229,328
Cost of identifying seropositive case — \$	40.71
Evaluation	
PCR assay for HCV RNA — \$†	14,981,084
Clinical, laboratory, and ultrasonographic evaluation — \$‡	31,349,925
Total cost of evaluation — \$	46,331,009
No. with viremia	1,605,116
Cost of HCV RNA testing and evaluation per viremic case — \$	<u>28.86</u>
Cost of identifying viremic case — \$	<u>85.41</u>
Treatment — \$	
Total cost of treatment	70,041,432
Cost of treatment per case§	<u>43.64</u>
Total cost	
Total cost of screening, evaluation, and treatment — \$	<u>207,139,177</u>
Cure rate — %¶	98.8
Cost of identifying and curing a case — \$	<u>130.62</u>

* All costs are in U.S. dollars, calculated at the exchange rate at the start of the program in October 2018 (1 U.S. dollar = 17.6 Egyptian pounds). Costs of screening include all costs incurred to screen the 49.6 million persons for HCV antibodies. Costs of evaluation and treatment assume that all seropositive patients were evaluated and that all patients with viremia were treated. PCR denotes polymerase chain reaction.

† Included are the purchase cost per test (\$4.80) plus 40% overhead, consumables, and staff, multiplied by the number of HCV-seropositive cases identified in the screening program.

‡ Included are the cost of laboratory tests and ultrasonography, consumables, and staff, multiplied by the number of HCV-seropositive cases identified in the screening program.

§ Included is the cost of a 12-week supply of locally manufactured sofosbuvir plus daclatasvir with or without ribavirin.

¶ Shown is the percentage of patients with a known sustained virologic response in the program.

|| The result is the cost of screening, evaluation, and treatment for the whole program divided by the number of patients with viremia divided by the cure rate.

200 Millions USD pour 1 million de patients traités en deux années

50 millions à traiter = 10 milliards de dollars

Objectif > 2030

*I. Waked et al, Special report Screening and Treatment Program to Eliminate Hepatitis C in Egypt
n engl j med 382;12 nejm.org March 19, 2020*



CONCLUSION

- Le rendez vous de 2030 sera possible pour les pays à revenu élevé avec une prévalence acceptable de patients virémiques : 20 % d'élus .
- Bassin méditerranéen : 3 pays , France- Italie- Espagne
- Le scénario Égyptien est l'exemple à suivre pour les pays de la rive sud